

A note on regularization and renormalization

M. Stenmark^{*1}

¹*Department of Radiation Sciences
Uppsala University
Box 535, S-75121 Uppsala*

We look at sections of a function bundle over the space of linear differential operators. We find that one can construct an isomorphism between a certain quotient bundle and the fourier counterpart of the original bundle defined by formal integration by parts. We also show that differential renormalization is an example of this technique.

I. INTRODUCTION

The systematics of renormalization rests on solid ground since several years. The problem of singular functions that comes from an ill defined Lagrangian can be taken care of via the procedure of introducing counter terms. If in addition the Lagrangian is renomalizable it is enough to add one set of counterterms which takes care of the infinities to all orders[1]. In this paper we will consider certain function bundles whose sections may be "regularized" with the help of the linear differential operator in the space of linear operators which it is a section above. Furthermore the Fourier transform will prove to be a homeomorphism onto the Fourier counterpart of the bundle. In this bundle the functions will yield finite results. In the next section we will show that one may define a quotient bundle that is isomorphic to the Fourier bundle and study how this comes about. We study why the results become finite and why this is actually what happens in differential renormalization.

II. REGULARIZATION AND RENORMALIZATION OF THE FUNCTION BUNDLE

Consider the space of linear differential operators, call it ξ . Consider the function algebra consisting of functions with well defined Fourier transform together with the functions generated by applying an operator to these. The added functions do not necessarily have a Fourier transform. This algebra is for each operator $L \in \xi$ a section over the point $L \in \xi$, call it $\mathcal{A}_L$. The function bundle we will study is the one spanned by these sections.

First we define the Fourier transform of a function in $\mathcal{A}_L$. We do so by formal integration by parts where the surface terms are neglected. If the function is in the sub-algebra consisting of functions with Fourier transform $\mathcal{B}$ the surface terms are zero so the only change is for the functions generated by L . For such a function one has $f = Lg = \sum a_i \partial^i g$ and thus performing formal integration by parts results in a convolution of the Fourier

transform of derivatives of the operator coefficients a and the fourier transform of g . If the operator consists of only derivatives this gives just multiplication by a power of p . This map is a homeomorphism between $\mathcal{A}_L$ and $\hat{\mathcal{A}}_L$.

If we now construct an ideal $\mathcal{I}_L$ of the algebra we may take quotient with this ideal and construct a quotient algebra $\mathcal{A}_L/\mathcal{I}_L$. First we define a character $\epsilon : \mathcal{B} \rightarrow \mathcal{C}$ by $\epsilon(f \in \mathcal{B}) = \hat{f}(p)$ where we chose p in the character such that it coincides with the map $\mathcal{F}$. With this we may define an ideal of $\mathcal{A}$ as $\mathcal{I} = \{a(b - \epsilon(b)) | b \in \mathcal{B}, a \in \mathcal{A}\}$.

It is now strait forward to use the fundamental homeomorphism theorem [2] to show that the below diagram commutes.

$$\begin{array}{ccc} \mathcal{A} & \rightarrow & \hat{\mathcal{A}} \\ \downarrow & \nearrow & \\ \mathcal{A}/\mathcal{I} & & \end{array} \quad (1)$$

To prove this we observe that it is enough to show that $\mathcal{I}$ is in the kernel of the homomorphism between $\mathcal{A}$ and $\hat{\mathcal{A}}$. But due to the choice of p that coincides with $\mathcal{F}$ this is trivial. Thus the isomorphism between the quotient bundle and the fourier bundle is induced by the above diagram.

One may argue that the skipping of surface terms is unphysical, but it turns out that this can be seen as a counterterm procedure in the following way. Assume that the function is determined from some Lagrangian and depends only on the square of x_μ . And that L fulfills the same invariance. The surface terms in n dimensional euclidian space can be written,

$$\int_{S_\epsilon} d\sigma_\mu f(x) \left(\int dx_\mu Lg \right) = \epsilon^{n-1} \frac{2\pi^{n/2}}{\Gamma(n/2)} f(0) \left(\int_{x^2=\epsilon^2} dx_\mu Lg \right) + \mathcal{O}(\epsilon) \quad (2)$$

where S_ϵ is the surface of an epsilon ball about the origin. We assume above that Lg obeys euclidian invariance. This terms times a constant can be added to the Lagrangian as a counterterm as in [3] and the constant determined to cancel the surface term in $\mathcal{F}$. Further trouble may still arise at the boundary of an infinite sphere but in most cases where f is an exponential this will be damped out by oscillations [3].

^{*}email: marten.stenmark@tsl.uu.se

The procedure of differential regularization and renormalization is an example of the above. Take ϕ^4 theory as an example and look to the point $(\partial^2, 1/x^4)$. In the function bundle one has a logarithm with a free mass parameter as the function which ∂^2 acts upon to produce $1/x^4$. The free parameter traces out a path through the section $\mathcal{A}_{\partial^2}$ of the function bundle. The Fourier map takes this path onto a path through the corresponding Fourier section.

Future work must look into the problem of symmetries, and among them those which generate Ward identities. Due to the fact that regularization may invalidate the Ward identities among greens functions these are only formal for bare Green functions [4]. Although the function bundle contains bare Green functions it is in no way trivial to show that the map induced by the diagram will obey these symmetries and thus generate appropriate Ward identities which hold globally on the bundle.

Symmetries such as gauge symmetries can be proved to be fulfilled on certain paths through the bundle [3, 4] but to see if some general results may be obtained further work is necessary. The cases which has been checked using DR[3] obeys Callan-Symanzik equations for the free parameters which trace out the path through the sections of the bundle. Generally one will have different integration constants for each $L \in \xi$. These can be coupled to each other using modified Ward identities that couple different paths together. So far this has just been done for a few examples mentioned earlier.

The bundles presented above gives a new way of viewing renormalization since we can look at the bundle of renormalizable functions as a whole. Certainly much work lie ahead if one is to understand the total structure of this construction, but the pavement is made and thus one may start the process of generalizing the ideas which has been set on formal basis during the past decades.

-
- [1] G. 't Hooft and M. J. Veltman, Nucl. Phys. B **44**, 189 (1972).
 - [2] A. I. Kostrikin and I. R. Shafarevich, Encyclopedia of Mathematical Sciences 11. Springer Verlag(1990).
 - [3] D. Z. Freedman, K. Johnson and J. I. Latorre, Nucl. Phys.

- B **371**, 353 (1992).
- [4] F. del Aguila, A. Culatti, R. Munoz Tapia and M. Perez-Victoria, Nucl. Phys. B **537**, 561 (1999) [arXiv:hep-ph/9806451].